

The medullary thick limb: Function and modulation of the single-effect multiplier

STEVEN C. HEBERT, W. BRIAN REEVES, DONALD A. MOLONY, and THOMAS E. ANDREOLI

Division of Nephrology, Department of Internal Medicine, University of Texas Medical School at Houston, Houston, Texas, USA

The specialized transport properties of the thick ascending limb of Henle's loop (TALH) enable this segment of the nephron to dissociate salt and water absorption. Sodium chloride absorption by the cortical (cTALH) and medullary (mTALH) thick ascending limb accounts for urinary dilution, since each of these segments is virtually water-impermeable. Salt absorption by the mTALH also contributes to medullary interstitial hypertonicity, supplies the energy for the "single effect" of countercurrent multiplication, and hence modulates urinary concentrating mechanisms. The medullary (mTALH) and cortical (cTALH) portions of the TALH are clearly heterogeneous, and can be distinguished on the basis of morphological [1, 2], biochemical [3–6] and functional [7] differences.

This article will focus particularly on the transport functions of the medullary segment. The specific issues discussed will include the mechanism of NaCl absorption by the mTALH, the enhancement of this salt absorptive process by antidiuretic hormone (ADH), at least in certain mammalian species, and the major factors that appear to regulate or modulate both ADH-dependent and -independent NaCl transport processes. Sufficient information is currently available which indicates that these modulating factors provide feedback loops which sense interstitial osmolality and serve as a means of controlling the magnitude of interstitial hypertonicity, and hence concentrating power.

Some transport properties

In vivo micropuncture studies indicated that dilution of the urine emerging from the loop of Henle resulted from NaCl abstraction from the thick ascending limb of Henle [8]. In vitro micropuncture studies on the mTALH [9, 10] demonstrated four cardinal features of NaCl absorption by the medullary segment. First, Cl^- absorption occurred against a transepithelial electrochemical gradient. Second, net salt absorption resulted in a relatively large lumen-positive transepithelial voltage (V_e , mV) which was sensitive to the loop diuretic, furosemide, when applied from the luminal side. Third, both net Cl^- absorption and the transport-related transepithelial voltage depended on the activity of $(\text{Na}^+ + \text{K}^+)\text{-ATPase}$, present in large amounts along the basolateral membrane of this nephron segment [3]. Finally, the transepithelial ion conductance of the

mTALH, calculated from passive tracer permeability measurements, indicates that this nephron segment is electrically "leaky" and that the shunt pathway is predominantly cation selective [7, 9, 10]. This latter feature is particularly unusual among epithelia with negligibly low water permeabilities.

Table 1 presents a summary of the important transport properties of the rabbit, rat and mouse mTALH that are relevant to the concentrating and diluting functions of this nephron segment. While it is clear that the mTALH segments from these species are quite similar with regard to the features discussed above, there are two significant differences between the rat (mouse) and rabbit. As these differences may play important roles in the concentrating process, it should also be noted that the maximal concentrating abilities of the rat and mouse are much greater than that of the rabbit.

The first important difference in the functional properties of the mTALH among these species is with regard to their urea permeabilities (P_{urea}). Knepper [11] has recently observed that the urea permeability of the rat mTALH is an order of magnitude greater than that of the rabbit (Table 1). Consequently, he has proposed that the P_{urea} of the TALH from the outer stripe of the rat outer medulla may be sufficient to permit the passive absorption of urea from this segment. This process not only would contribute to urinary dilution, by solute removal, but might also provide a pathway for medullary urea recycling, and thereby promote urinary concentration. Comparable data are not available on the urea permeability of the mTALH from mice.

The second major functional difference among these species is the much greater ADH-responsiveness of the adenylate cyclase system in rat and mouse mTALH segments [4–6]. The enhanced ability of ADH to stimulate this enzyme activity also appears to correlate with differences in the maximal concentrating abilities of the rat (mouse) and rabbit.

Site limiting mTALH water transport

The major functional property of the mTALH that permits this segment to dilute the urine is its negligible transepithelial water permeability. Since the ratio of junctional interspace to apical cell membrane surface area for the TALH is quite low [12], it seems reasonable to argue that the transcellular water permeability of this segment also be quite low. Recent qualitative and quantitative morphological observations in the mouse mTALH have indicated that the major barrier limiting transcel-

Table 1. General characteristics of medullary thick ascending limbs

Species	ADH	P_f	$P_{D_{urea}}$	P_{Na}	P_{Cl}	$(Na^+ + K^+)-ATPase$ $\mu moles\ cm^{-1}\ sec^{-1}$	Adenylate cyclase
		$\mu m\ sec^{-1}$					
Mouse	-	6-23		0.23	0.10		
	+	6-23		0.25	0.12	10	++
Rabbit	-	<5	<0.1	0.63	0.11		
	+	<5				7	++
Rat	-	≈ 0	0.7(IS), 1.4(OS)				-
	+	≈ 0				13	+

The data are from Refs. 3, 5, 6, 7, 10, 13.

lular water movement in this nephron segment resides at the apical (luminal) membrane [13].

Figure 1 shows the effects of perfusate and peritubular bath dilution on the tubular cell morphology of a single in vitro perfused mouse mTALH [13]. ADH was present throughout the experiment at a concentration that maximally stimulates NaCl absorption in this segment. During the initial period, isotonic Krebs-Ringer bicarbonate buffer (290 mOsm/liter; KRB) was present in both perfusing and bathing solutions. No change in the tubular cell morphology or estimated cell volume [13] was observed during about a six minute period after changing the perfusate to the dilute solution. However, very rapidly following bath dilution, tubular cells were seen bulging into the lumen (Fig. 1) and cell volume increased by 30%. These observations indicate that apical cell membranes of the mouse mTALH are virtually water-impermeant even in the presence of ADH. Therefore this membrane limits transepithelial water absorption in this region of the diluting segment.

General model of NaCl absorption

Analyses [7, 14-18] of the specific transport mechanisms involved in NaCl absorption by the mTALH, and the origin of the lumen-positive transepithelial voltage, may be integrated into the model shown in Figure 2. This model shares many similar features with those proposed for the rabbit cTALH [19] and non-mammalian diluting segments [20, 21]. The specific data in support of the model shown in Figure 2 have been reviewed elsewhere [22, 23]; we will therefore focus on some of the general features of the model.

Net Cl^- absorption by the mTALH involves a secondary active transport process in which luminal Cl^- entry into cells is mediated by an electroneutral $Na^+K^+:2Cl^-$ co-transport process that is inhibited specifically by loop, or "high ceiling," diuretics. Recent studies assessing the ion dependence of Cl^- absorption by the in vitro perfused mouse mTALH indicate clearly that net Cl^- absorption requires both Na^+ and K^+ in luminal fluids [7, 15]. Measurements of tracer Cl^- uptake by either membrane vesicles or cells isolated from the medulla have [24-26], with few exceptions [27, 28], also demonstrated a dependence on both external sodium and potassium. In a recent report, Eveloff, Fong and Calamia [29] have indicated that a K^+ dependence for tracer Cl^- uptake in isolated rabbit mTALH cells could be demonstrated only after subjecting cells to hypertonic media. However, it is not clear whether this represents an intrinsic property of mTALH cells, or whether it is related to the method of preparation of the cells. Further evidence for the interdependence of Na^+ , K^+ and Cl^- in the

mTALH has been provided by Forbush and Palfrey [30], who demonstrated that the binding of (3H)-bumetanide by apical membrane vesicles prepared from the outer medulla of dog kidneys was dependent on the presence of all three ions.

No direct measurements of the electrochemical gradients for co-transport of all three ions across the apical membrane of the mammalian mTALH have yet been made because of the difficulty in determining intracellular ion activity in these small cells. However, a favorable integrated electrochemical gradient for the electroneutral co-transport of $Na^+K^+:2Cl^-$ units has been established for the amphibian diluting segment which possesses a similar mechanism for NaCl absorption [20, 31, 32]. At the present time, it seems reasonable to argue that the electrochemical gradient for Na^+ across the apical membrane is sufficient to drive the uphill movement of K^+ and Cl^- into cells. Thus apical Cl^- entry into cells depends ultimately on basolateral membrane $(Na^+ + K^+)-ATPase$. Consequently, ouabain addition to, or K^+ removal from, peritubular fluid inhibits salt transport in the mTALH [7, 10].

The exit of Cl^- from cells across the basolateral membrane is primarily conductive [15, 16, 18], although a small component of electroneutral KCl co-transport, which has been suggested for the rabbit cTALH [33], cannot be excluded entirely for the mTALH. The large intracellular negative voltage in the mouse mTALH [16] provides a portion of the driving force for electrical Cl^- exit. Based on the electrical properties of basolateral cell membranes, we have estimated [34] the intracellular Cl^- activities required to account for observed rates of Cl^- absorption: without ADH, the calculated Cl^- activity is 16 mM; and with ADH, 25 mM. Similar values for intracellular Cl^- activity have been measured in the rabbit cTALH without ADH [35].

An important characteristic of the mTALH, as well as all diluting segments, is the presence of a Ba^{++} -sensitive apical membrane K^+ conductance [15-17, 35]. The apical K^+ channels in the mouse mTALH share some of the characteristics of Ba^{++}/K^+ interactions observed in excitable tissue (such as, both voltage- and concentration-dependence and Ba^{++}/K^+ competitive effects [15-18]). Furthermore, these K^+ channels represent the sole conductive pathway identified to date in apical membranes of the mouse mTALH [15-18]. A role for this K^+ conductance in NaCl absorption by the mouse mTALH is indicated by the observation that luminal Ba^{++} can effectively abolish net Cl^- absorption [15-17, 34], presumably because luminal Ba^{++} depolarizes the cell and reduces conductive Cl^- exit across basolateral membranes [15, 16].

Recent estimates of the magnitude of the paracellular (and lateral intercellular space) conductance have provided a clearer




Fig. 1. The rate limiting for transcellular water permeability in the mouse *mTALH*. A single mouse *mTALH* segment was perfused in vitro at 37°C in the presence of bath ADH at 25 μ U/nl. The perfusate and bath contained Krebs-Ringer's bicarbonate (KRB) media, 290 mOsm/KgH₂O. The hypotonic KRB media were produced by reducing the NaCl concentration by 25 mm/liter: final osmolality was 240 mOsm/KgH₂O.

picture of the important role played by both the apical K⁺ conductance and the cation-selective paracellular pathway in transepithelial NaCl absorption by the mouse *mTALH* [17]. High concentrations of luminal Ba⁺⁺ (20 mM, at zero luminal K⁺) were used to block transcellular conductance, so that the residual transepithelial conductance represented, virtually exclusively, the paracellular pathway. Then a large reversed (lumen-to-bath, as opposed to normal luminal dilution) osmotic gradient was applied across the tubule by adding large concen-




Fig. 2. General model for NaCl absorption in the mouse *mTALH*. From [15–17, 23].

trations of urea to luminal fluids. This maneuver produced a concentration-dependent increase in electrical conductance and a reduction in the Na⁺/Cl⁻ permselectivity ratio of the paracellular pathway to free solution values. These functional changes were accompanied by blistering at the luminal surface similar to that found with apical hypertonicity effects in other water-tight epithelia [36].

These osmotic effects were referable to opening of the junctional complexes; thus the residual transepithelial conductance measured with luminal 20 mM Ba⁺⁺ and with luminal 900 mM urea provided a minimal estimate of the lateral intercellular space resistance. This latter value was then used to calculate a maximal estimate of the concentration of NaCl in lateral interspace fluid that would occur during transepithelial NaCl absorption. The contribution of interspace NaCl accumulation to the transepithelial voltage could be determined from these data.

The results of these studies [17] indicate that the paracellular spaces are in virtual diffusion equilibrium with bulk solutions, so that transcellular, rather than paracellular, transport events must account for the origin of the transepithelial voltage. In other words, the differing ionic conductance properties of apical (principally K⁺) and basolateral (primarily Cl⁻) membranes appear to provide the critical pathways for the development of the lumen-positive transepithelial voltage. These observations, together with the close correspondence between the measured equivalent short circuit current and the sum of net Cl⁻ and K⁺ transport in the mouse *mTALH*, indicate that Cl⁻ absorption is rheogenic, that is, associated with transcellular and paracellular current flows. According to this rheogenic model, the K⁺ that enters the cell via the apical electroneutral salt co-transporter recycles, in large part, conductively across apical membranes, thus providing the apical component of the transepithelial current. In other words, K⁺ recycling across apical membranes provides a means for maintaining electroneutral Na⁺:K⁺:2Cl⁻ entry across apical membranes, and blockade of the apical K⁺

Table 2. Transport properties of mouse mTALH segments

V_e mV	J_{Cl}^{net} $pEq\ sec^{-1}\ cm^{-2}$	Shunt			$J_{Cl}^{net}:shunt\ J_{Na}^{net}$ ratio
		P_{Na}/P_{Cl}	G_{Na} ($mS\ cm^{-2}$)	J_{Na}^{net} ($pEq\ sec^{-1}\ cm^{-2}$)	
9.7 ± 0.5	$10,950 \pm 1180$	3.0 ± 0.3	45.5 ± 4.7	4530 ± 550	2.4 ± 0.3

Data obtained from 13 mouse mTALH segments perfused in vitro [17] with ADH $10\ \mu U/ml$ present in the peritubular bath throughout. Values represent mean \pm SEM. The shunt values were calculated as indicated below while the tubules were perfused with $20\ mM\ Ba^{++}/zero\ K^{+}$ so that the transcellular conductance pathway was completely blocked. The paracellular Na^{+} to K^{+} permselectivity ratio (P_{Na}/P_{Cl}) was calculated from dilution voltage observed during reductions in bath $NaCl$ concentration. The shunt $J_{Na}^{net} = 1/[F/G_s\ V_e\ (1 + 1/(P_{Na}/P_{Cl}))]$, where G_s is the residual transepithelial conductance in the presence of $20\ mM\ Ba^{++}/zero\ K^{+}$ and F is the Faraday. The shunt $G_{Na} = 1/[G_s\ (1 + 1/(P_{Na}/P_{Cl}))]$. The net transcellular Cl^{-} absorption (J_{Cl}^{net}) was calculated as $J_e - J_{K}^{sec} = J_e - 1/[F/G_s\ V_e\ (1 + P_{Na}/P_{Cl})]$, where J_e is the equivalent electrical flux (I_{sc}/F).

conductance pathway by lumen Ba^{++} abolishes the transepithelial voltage because transepithelial currents are stopped.

One other prediction of the model shown in Figure 2 is also supported by these results. Since dilution of the urine by the mTALH occurs through the net absorption of equal quantities of Na^{+} and Cl^{-} , a stoichiometry of $1Na^{+}:2Cl^{-}$ for the transcellular salt transport mechanism requires that one-half of the net Na^{+} absorbed flows via the paracellular route. We have recently calculated the ratio of measured net Cl^{-} absorbed transcellularly to the net Na^{+} that could be absorbed through the paracellular route [17]. The latter net Na^{+} flux was estimated from the measured paracellular conductance, the electrical P_{Na}/P_{Cl} ratio and the observed magnitude of the transepithelial voltage. The results of these measurements are shown in Table 2. In accord with the prediction of the model shown in Figure 2, the magnitude of the transepithelial voltage in each tubule provides a driving force sufficient to transport a quantity of Na^{+} through the cation-permeable pathway approximately equal to one-half the rate of net Cl^{-} absorption.

The mechanism of $NaCl$ absorption in the mTALH shown in Figure 2 has at least two general consequences. First, rheogenic Cl^{-} absorption in the mTALH may reduce the energy expenditure required for transepithelial Na^{+} transport. As discussed above, the combination of a lumen-positive transepithelial voltage and a high conductance, Na^{+} -selective shunt pathway permits the transport of about 50% of the total Na^{+} absorbed through the paracellular route. Thus compared to epithelia that transport all the Na^{+} absorbed through the transcellular route, the O_2 utilized to absorb a fixed quantity of Na^{+} would be reduced to one-half. That is, assuming a fixed $3Na^{+}:2K^{+}$ stoichiometry for basolateral membrane ($Na^{+} + K^{+}$)-ATPase, about 18 moles of Na^{+} are transported transcellularly per mole of O_2 consumed; this may increase to approximately 36:1 for the mTALH [23].

Since the mTALH is generally thought to be in an in vivo environment with a low O_2 tension [37], the greater efficiency of its salt transport mechanism may be important for metabolic survival. In fact, recent observations in the isolated perfused rat kidney by Alcorn, Emslie and Boss and by Brezis et al [39] have demonstrated that the mTALH is uniquely sensitive to metabolic damage, due either to reduced O_2 delivery or to increases in transcellular ion transport. This phenomenon may account for the early loss of concentrating ability in certain forms of acute renal failure.

There may be one other advantage to the salt transport mechanism depicted in Figure 2. Schultz [40] has stressed the importance of homoregulatory processes in salt transporting

Table 3. Effect of ADH on $NaCl$ and water transport in the mTALH

Species	ADH	G_e $mS\ cm^{-2}$	J_{NaCl} $pM\ sec^{-1}\ cm^{-2}$	V_e mV	References
Mouse	—	80	2600	5	[7, 15, 43, 44]
	+	100	10,800	11	[7, 15, 43, 44]
Rabbit	—		6400	3–7	[10, 44]
	+		unchanged	unchanged	[10]
Rat	—		3450 ^a	2.5	[44]
	+		9070 ^a	3.6	[44]

^a Calculated from [44] assuming an inner tubule diameter of $20\ \mu m$.

epithelia which adjust the rate of Na^{+} entry into cells to equal the rate of Na^{+} exit, in order to avoid potentially lethal changes in cell volume. These processes are particularly important in epithelia like the mTALH where the rate of transepithelial salt transport can change rapidly. A mechanism that permits one-half of the net Na^{+} movement to proceed via the paracellular route—and not affect cell volume—might be beneficial in the mTALH whose rate of $NaCl$ absorption is about equal to, but whose cell volume is only about one-third that of the proximal tubule.

Regulation of $NaCl$ absorption

Stimulation of $NaCl$ absorption by ADH

Wirz [41] first suggested that ADH might regulate counter-current multiplication by augmenting the rate of $NaCl$ abstracted from the ascending limb of Henle's loop. Added support for this possibility obtained by the recent observations of ADH-induced increases in adenylate cyclase [4–6] and protein kinase [42] activities in medullary thick ascending limbs of rabbit, rat and mouse.

Table 3 summarizes the effects of ADH on salt transport in the in vitro perfused mTALH segments of mouse, rat and rabbit. Hall and Varney [43] established that ADH increased the lumen-positive transepithelial voltage and the net rate of tracer Cl^{-} absorption in the mouse mTALH. Sasaki and Imai [44] and we [7, 14] confirmed these effects of ADH (and cyclic AMP) on salt transport in this species; this hormone effect was restricted to the medullary segment of the TALH [7, 14]. Moreover, the maximal effect of ADH on salt transport occurred at hormone concentrations found during antidiuretic states [7, 45]. Likewise, in homozygous Brattleboro rats, ADH (or 1-desamino-8-D-arginine vasopressin [dDAVP]) increases $NaCl$ absorption in the mTALH of this species when assessed

by in vitro or by in vivo microperfusion of the loop of Henle [46, 47]. In contrast, no effect of pharmacologic concentrations of ADH (or cyclic AMP) could be demonstrated on salt transport in the rabbit mTALH [44] despite the significant, albeit variable, stimulation of adenylate cyclase by hormone in this species [4–6]. Insight into the varied effects of ADH on salt transport by the mTALH of these species may be gained by considering the interactions of ADH and osmolality. Since the osmotic environment of the mTALH varies with the state of water balance, such osmotic interactions may be quite important to hormone action in vivo. In renal medullary tissue, the effects of osmolality on both basal and ADH-stimulated adenylate cyclase activity are complex [48–52]. Increasing the NaCl concentration has a biphasic effect, with concentrations below 200 mM stimulating, but greater concentrations inhibiting, adenylate cyclase activity. However, urea consistently inhibits this enzyme at all concentrations above 50 mM. In contrast, DeRubertis and Craven [50] have observed vasopressin-stimulated cyclic AMP generation in renal medullary tissue increases in external osmolality up to 2000 mOsm/kg.

Torikai and Imai [52] have recently assessed the effects of osmolality on vasopressin-stimulated cyclic AMP generation in isolated fragments of rat mTALH. They find that hormone-stimulated cyclic AMP generation increases as a function of external NaCl, but not urea, concentrations. These results are consistent with the observations of Sasaki and Imai [44] that ADH stimulated NaCl transport in the isolated perfused rat mTALH only when the external fluids were hypertonic. Whether such osmolality effects can account for the apparent lack of hormone stimulating salt transport in the rabbit mTALH is unknown.

The issue of whether ADH can stimulate NaCl transport in the human mTALH is not clear. Chabardés and co-workers [4] were unable to detect any effect of ADH on cyclic AMP production in human mTALH segments isolated from pump-perfused kidneys. However, Alcorn, Emslie and Boss [38] have shown that, in isolated perfused rat kidneys, marked morphological changes occur in the mTALH within five minutes after starting perfusion. Given the unique sensitivity of the mTALH to hypoxic damage [39], it is possible that, in the pump-perfused human kidneys, hormone effects could be blunted.

ADH-prostaglandin interactions

The metabolism of renal medullary prostaglandins, and the interactions between ADH and these lipids, have been the subject of several recent reviews [53–55]. Prostaglandins are synthesized and rapidly inactivated in the renal medulla; their primary function is local. Prostaglandin E₂ is the major product of prostaglandin synthesis in the renal medulla, and is responsible for the major physiological effects of prostaglandins on water excretion.

At least two lines of argument suggest that PGE₂ may participate in a medullary feedback loop that modulates the rate of NaCl absorption by mTALH. First, PGE₂ has been shown to reduce directly the rate of net NaCl absorption in the mTALH, but not in the cTALH [56, 57]. Likewise, the possibility that PGE₂ inhibits in vivo NaCl absorption by the mTALH is supported by the micropuncture observations of an increase in ²²Na⁺ excretion in the urine when tracer ²²Na⁺ and PGE₂ were injected simultaneously into the late proximal tubule of the rat

[58]; by the finding of a rise in NaCl content of the medulla when prostaglandin synthesis is inhibited [59]; and by alterations in loop chloride transport when inhibitors of prostaglandin synthesis are given [60, 61]. In contrast, Stokes [62] found that PGE₂, in micromolar concentrations, enhanced the ADH-unstimulated transepithelial voltage and net Cl[−] absorption by the rabbit mTALH; however, an inhibiting effect of rabbit serum may have obscured a separate suppressive effect of PGE₂ on NaCl absorption [55, 57].

Second, there are specific interactions between PGE₂ and ADH in regulating NaCl absorption by the in vitro microperfused mouse mTALH [56, 63]. In this species, as opposed to the rabbit, micromolar concentrations of PGE₂ have no effect on NaCl absorption or the transepithelial voltage in the absence of ADH. However, in the presence of ADH, that is, when the rate of NaCl absorption is maximal, PGE₂ reduces the ADH-dependent component of NaCl transport in a dose-dependent fashion with half maximal inhibition occurring at about 10^{−10} M PGE₂. Likewise, PGE₂ inhibits ADH-dependent, but not basal, cellular AMP accumulation in isolated rat mTALH segments [64].

The PGE₂-ADH interactions in the mTALH are seemingly competitive, since supramaximal concentrations of ADH can reverse the inhibiting effect of PGE₂ on NaCl absorption [56, 63]. It now appears probable that PGE₂ inhibits the action of ADH by stimulating an inhibitory guanine nucleotide regulatory subunit of the adenylate cyclase system [63]. A similar mechanism may apply to PGE₂-ADH interactions on water permeability in the collecting duct [64]. A similar interaction likely occurs between PGE₂ and ADH in the rat.

It is unlikely that PGE₂ production by mTALH cells contributes to in vivo ADH-PGE₂ interactions, since cyclooxygenase activity is negligibly small in this nephron segment [66]. However, PGE₂ production may occur in other medullary components, including the medullary collecting duct and interstitial cells. PGE₂ synthesis by medullary/papillary interstitial cells can be modulated both by ADH and external solution osmolality [67]. These modulating factors appear to work by altering the calcium-dependent acyl hydrolase activity that regulates the availability of arachadonic acid for PGE₂ production by these cells.

The interaction among these modulating factors is complex. ADH can directly and acutely stimulate the synthesis of PGE₂ by interstitial cells, an effect that seems related primarily to the pressor, rather than the antidiuretic, activity of ADH [53, 54]. In contrast, during chronic ADH administration, antidiuretic activity may be important to an increase in cyclooxygenase activity [53, 54]. The role of ADH-PGE₂ interactions during physiological antidiuresis is unclear, since water deprivation in man [68] and animals [69, 70] may be associated with reductions in urinary PGE₂ excretion. Hypertonic NaCl also stimulates the release of PGE₂ from interstitial cells. Both this stimulation and that due to ADH can be reduced or abolished by hypertonic urea [67].

Osmolality effects on NaCl absorption

Medullary interstitial osmolality might regulate the rate of net NaCl absorption by the mTALH in at least two ways. First, increases in interstitial NaCl concentration produced by active NaCl absorption from the mTALH increase the driving force

for passive salt leakage from interstitium to luminal fluid via the paracellular pathway. Second, peritubular hypertonicity may down-regulate directly net salt absorption. Thus in the *in vitro* microperfused mouse mTALH, increases in the osmolality of peritubular bathing solutions with urea rapidly and reversibly inhibit the rate of active NaCl absorption [71]. This effect is not referable to the transtubular osmotic gradient, but rather to the magnitude of the increase in peritubular osmolality [71]. Since hypertonic urea does not affect either basal or ADH-stimulated cyclic AMP generation in the rat mTALH [52], it seems likely that processes beyond the generation of cyclic AMP participate in this urea-mediated inhibition. This view is supported by the observations that neither supramaximal concentrations of ADH nor dibutyryl-cyclic AMP are able to reverse the inhibition of NaCl transport mediated by hypertonic peritubular bath urea [71].

Other regulators of NaCl absorption

Adrenergic agonists. Adrenergic agonists can promote an effect similar to that produced by ADH in both man [72] and animals [73, 74]. Isoproterenol stimulates NaCl absorption by the mouse mTALH, and this effect can be inhibited by propranolol [75, 76]. Glucagon also augments the spontaneous trans-epithelial voltage in the *in vitro* microperfused mouse mTALH [76]. While these observations suggest that multiple agents may interact to regulate NaCl absorption by the mTALH [77], it should be noted that both isoproterenol and glucagon have affinity constants for action on the mTALH some three orders of magnitude higher than that for ADH, and that neither agent is additive to the effect of ADH [76].

Barajas and coworkers [78, 79] have demonstrated the presence of adrenergic nerve terminals along the basolateral membranes of the cTALH and mTALH in the rat. Furthermore, DiBona and Sawin [80] have shown that low frequency renal nerve stimulation increases the rate of salt absorption from the loop of Henle in both hydropenic and isotonic volume expanded rats; and, Bencs  th, Sz  n  si and Tak  cs [81] have observed reductions in loop NaCl absorption acutely following renal sympathectomy. These latter observations are consistent with the possibility that *in vivo* adrenergic nerve stimulation modulates salt transport by the TALH.

Medullary K^+ concentrations. Increased medullary K^+ concentrations occur by medullary K^+ recycling [82, 83]. The mTALH may play an important role in the recycling process [83, 84], and vasopressin may also stimulate medullary K^+ recycling by an action on the mTALH and on K^+ secretion in the distal tubule [15, 16, 85, 86]. The magnitude of medullary K^+ concentrations in the medulla are thought to be important in renal K^+ excretion [82–84]. Stokes [84], for example, observed a decrease in the rate of NaCl absorption from the *in vitro* microperfused mTALH following an increase in the external K^+ concentration from 5 to 25 mM, and suggested that the resultant increase in NaCl delivery to the distal convoluted tubule and collecting duct may enhance renal K^+ excretion. According to the model for salt absorption in the mTALH shown in Figure 2, the increased external K^+ (either luminal or peritubular) could depolarize mTALH cells via K^+ conductive pathways in apical or basolateral membranes, and thereby reduce the driving force for conductive Cl^- exit across the




Fig. 3. General model for regulation of urinary concentrating processes. From [23].

basolateral membrane. An acute effect of K^+ loading on loop NaCl absorption has also been observed *in vivo* [86].

Potassium deprivation is also associated with reductions in urinary concentrating ability. The micropuncture studies of Luke, Booker and Galla [87] in K^+ depleted rats suggest that a reduction in loop NaCl absorption could, at least in part, account for the impairment. The reduced loop NaCl absorption could result either from a reduction in ADH-sensitive adenylate cyclase activity or from a direct effect on low external K^+ on NaCl absorption by the TALH. Kim et al [88] have recently demonstrated that vasopressin-dependent adenylate cyclase activity, but not cell cyclic AMP accumulation, was reduced in K^+ -depleted rats.

Calcium effects. Hypercalcemia reduces the ability to produce a maximally concentrated urine. This effect may be due, at least to some extent, to an inhibitory effect on ADH action [89, 90]. Since Ca^{++} may influence the adenylate cyclase–cyclic AMP system, it is possible that hypercalcemia may affect ADH-dependent cyclic AMP accumulation in the mTALH, a view consistent with the decreased medullary hypertonicity observed in the hypercalcemic state [89, 90]. Kusano et al [91] have provided evidence for an inhibitory effect of calcium on cyclic AMP accumulation in mTALH segments isolated from the rat. This effect appears to be due to a stimulation of phosphodiesterase activity. Thus, at least part of the hypercalcemic effect on urinary concentration may be by inhibition of ADH-stimulated NaCl absorption in the mTALH.

Model for integration of ADH, PGE_2 and osmolality effects

Figure 3 provides a model for integrating the effects of some modulators of urinary concentrating ability [23]. According to

the model shown in Figure 3, ADH-stimulated NaCl absorption by the mTALH is regulated by two negative feedback loops that "sense" the level of interstitial osmolality. In this manner, these feedback loops provide a means for controlling the magnitude of the driving force for water abstraction from collecting ducts, and thus urine concentration. During the early stage of antidiuresis, an ADH-mediated increase in NaCl absorption by the mTALH produces a rise in interstitial salt concentration [92–94]. The rise in interstitial NaCl concentration stimulates PGE₂ release from interstitial cells which, in turn, decreases the rate of ADH-dependent NaCl absorption. Since this ADH-PGE₂ interaction is concentration-dependent, alterations in ADH concentrations can influence the level of feedback. During antidiuresis, a rise in interstitial urea concentration decreases the hypertonic NaCl-mediated release of PGE₂ from medullary interstitial cells and directly inhibits salt absorption from the mTALH. It seems possible that this may provide a means of reducing the metabolic expenditure of energy by the mTALH in late antidiuresis, when high interstitial urea concentrations provide a major fraction of the driving force for water absorption from collecting ducts.

The ADH effect

It is now established that, in the mouse mTALH, ADH increases the rate of net salt absorption and the spontaneous transepithelial voltage V_e (Table 3). ADH also increases the transepithelial electrical conductance (G_e , mS cm⁻²) exclusively because of an increase in transcellular electrical conductance (G_c , mS cm⁻²); the hormone does not affect paracellular electrical conductance (G_s , mS cm⁻²) [34]. Finally, ADH augments the net rate of K⁺ secretion in mouse mTALH segments [15].

Some electrophysiologic considerations

In order to consider the mechanism for ADH action, it is pertinent to reiterate certain of the ionic conductance characteristics of these mTALH segments (Fig. 2). The apical membranes of the mouse mTALH are predominantly, and possibly exclusively, K⁺ selective; these apical K⁺ channels can be blocked by varying concentrations of luminal Ba⁺⁺, with zero luminal K⁺ [15–18, 34]. The blockade of apical K⁺ channels by Ba⁺⁺ is readily reversible, either by removing luminal Ba⁺⁺ and adding luminal 5 mM K⁺; or, in the presence of luminal Ba⁺⁺, by raising luminal K⁺ concentrations [15]. Cell impalement studies indicate that, with luminal Ba⁺⁺, the apical-to-basolateral membranes resistance ratio (R_a/R_b) rises about thirteenfold [16].

Finally, it is pertinent to note that with 20 mM luminal Ba⁺⁺ there is virtually complete, but readily reversible, blockade of apical K⁺ channels; and that the transcellular conductance G_c , when estimated in this manner, comprises approximately 40 to 50% of the total transepithelial conductance G_e [17, 34]. Consequently, in a given tubule segment one can measure: the transepithelial conductance G_e with luminal 5 mM K⁺, zero Ba⁺⁺; the shunt conductance G_s (mS cm⁻²) as the conductance insensitive to luminal 20 mM Ba⁺⁺, zero K⁺; and the transcellular conductance G_c as ($G_e - G_s$) [17, 34].

As noted in Figure 2, basolateral membranes contain (Na⁺ + K⁺)-ATPase, a K⁺ conductance, and a Cl⁻ conductance [15]. Net Cl⁻ absorption in the mTALH is virtually entirely conduc-

tive, and involves the basolateral Cl⁻ conductance [15, 17]. In agreement with this view, net Cl⁻ absorption may be abolished by applying Cl⁻ channel blocking agents to basolateral membranes [95].

An admittance mechanism for ADH

ADH may increase the net rate of salt absorption in the mTALH by way of a hormone-dependent increase in the functional number of Na⁺:K⁺:2Cl⁻ co-transport units in apical membranes [15, 16, 34]. Or put differently, ADH, acting through the second messenger cAMP, may exert an admittance effect on apical membranes. This postulate derives largely from an analysis of the interplay between ADH and furosemide in modulating transcellular ionic conductances. This argument, presented in detail elsewhere [34], may be summarized as follows.

Without luminal furosemide, ADH increases both the transcellular conductance G_c and the R_a/R_b ratio; thus the ADH-dependent increase in G_c involves an increase in the conductance of basolateral membranes. Moreover, without furosemide, the hormone-dependent increase in V_e occurs because V_{bl} , the basolateral membrane voltage, is depolarized to a greater extent than V_a , the apical membrane voltage. However, when luminal furosemide is present, ADH produces no detectable increase in G_c ; the R_a/R_b ratio is about 0.5–0.6, and unaffected by ADH; and V_a and V_{bl} are hyperpolarized to the same extent, so that V_e is virtually zero with or without ADH.

Thus when furosemide is used to block hormone-dependent increases in the rate of apical Na⁺:K⁺:2Cl⁻ entry—which must attend hormone-dependent increases in the rate of net salt absorption—there is no ADH-dependent increase in G_c . Put differently, these data are consistent with the view that ADH increases the functional number of apical membrane Na⁺:K⁺:2Cl⁻ co-transport units, and that G_c rises because the attendant changes in intracellular ionic activities result in a depolarization of V_{bl} and an increase in basolateral conductance [15, 34]. More specifically, it is plausible to argue that the latter increases because the ADH-dependent increase in apical Na⁺:K⁺:2Cl⁻ units raises intracellular Cl⁻ activity, and thus depolarizes V_{bl} . According to this view, basolateral conductance rises because of a voltage-dependent rise in the Cl⁻ conductance of basolateral membranes [34].

In this regard, removal of bath Cl⁻ produces a striking reduction in G_c , and this reduction is considerably greater with ADH than without ADH; in other words, ADH increases basolateral Cl⁻ conductance [34]. However, this ADH-dependent increase in basal Cl⁻ conductance is blocked by luminal furosemide [34]. Moreover, by using the change in G_c , which occurs with bath Cl⁻ removal in combination with R_a/R_b data, it has been possible to estimate the effects of ADH and furosemide on intracellular Cl⁻ activity [34]. Those calculations indicate that, without furosemide, ADH increases intracellular Cl⁻ activity from 16.3 mM to 25 mM; and that with furosemide, intracellular Cl⁻ is approximately 10.3 mM with or without ADH. These calculations are therefore congruent with the view that an ADH-dependent rise in intracellular Cl⁻ activity depolarizes V_{bl} , and produces a voltage-dependent increase in basolateral membrane conductance [34]. Thus, basolateral Cl⁻

channels in the mTALH may have properties analogous to Goldman rectifiers.

It has also been proposed that ADH might increase the number of Cl^- channels in basolateral membranes [18]. In quantitative terms, this effect appears small with respect to the ADH effect on apical $\text{Na}^+:\text{K}^+:2\text{Cl}^-$ co-transport units.

Rather, the primary effect of ADH on mTALH segments, by way of the second messenger cAMP [14], is to increase net Cl^- absorption by increasing apical membrane $\text{Na}^+:\text{K}^+:2\text{Cl}^-$ admittance. Such an apical membrane admittance effect for ADH in the mTALH is analogous to that of ADH in other epithelia: in mammalian cortical collecting tubules and in amphibian epithelia, ADH or cAMP increases the functional number of apical membrane water [96–99], Na^+ [100] and K^+ channels [101]. In these latter ADH-sensitive epithelia, the hormone also increases the rate of fusion of sub-apical membrane vacuoles into apical intramembranous aggregates [102].

Acknowledgments

S.C. Hebert is an Established Investigator of the American Heart Association. His current address is: Renal Division, Brigham and Women's Hospital, Boston, Massachusetts 02115, USA. W.B. Reeves is a Clinician-Scientist of the American Heart Association.

Reprint requests to Thomas E. Andreoli, M.D., Department of Internal Medicine, 1.150 MSMB, P.O. Box 20708, Houston, Texas 77225, USA.

References

1. ALLEN T, TISHER CC: Morphology of the thick ascending limb of Henle. *Kidney Int* 9:8–22, 1976
2. KAISLING B, KRIZ W: *Advances in Anatomy, Embryology and Cell Biology*, vol. 56, 1978
3. KATZ AI, DOUCET A, MOREL F: Na-K-ATPase activity along the rabbit, rat and mouse nephron. *Am J Physiol* 237:F114–F120, 1979
4. CHABARDÉS, D., GAGNAN-BRUNETTE M, IMBERT-TEBOUL M: Adenylate cyclase responsiveness to hormones in various portions of the human nephron. *J Clin Invest* 65:439–448, 1980
5. MOREL F: Sites of hormone action in the mammalian nephron. *Am J Physiol* 240:F159–F164, 1981
6. MOREL F, IMBERT-TEBOUL M, CHABARDÉS, D. Distribution of hormone-dependent adenylate cyclase in the nephron and its physiological significance. *Ann Rev Physiol* 43:569–581, 1981
7. HEBERT SC, CULPEPPER RC, ANDREOLI TE: NaCl transport in mouse thick ascending limbs. I. Functional nephron heterogeneity and ADH-stimulated NaCl cotransport. *Am J Physiol* 241:F412–F431, 1981
8. GOTTSCHALK CW, MYLLE M: Micropuncture study of the mammalian urinary concentrating mechanism: Evidence for the counter-current hypothesis. *Am J Physiol* 196:927–936, 1973
9. BURG MB, GREEN N: Function of the thick ascending limb of Henle's loop. *Am J Physiol* 224:659–668, 1973
10. ROCHA AS, KOKKO JP: Sodium chloride and water transport in the medullary thick ascending limb of Henle. *J Clin Invest* 52:612–623, 1973
11. KNEPPER M: Urea transport in isolated thick ascending limbs and collecting ducts from rats. *Am J Physiol* 245:F634–F639, 1983
12. WELLING LW, WELLING DJ, HILL JJ: Shape of cells and intercellular channels in rabbit cortical thick ascending limb of Henle. *Kidney Int* 13:144–151, 1978
13. HEBERT SC: Hypertonic cell volume regulation in mouse thick ascending limbs. I. ADH-dependency and nephron heterogeneity. *Am J Physiol (Cell Physiol)*. (in press)
14. HEBERT SC, CULPEPPER RC, ANDREOLI TE: NaCl transport in mouse medullary thick ascending limbs. II. ADH enhancement of transcellular NaCl co-transport; origin of transepithelial voltage. *Am J Physiol* 241:F432–F442, 1981
15. HEBERT SC, FRIEDMAN PA, ANDREOLI TE: Effects of antidiuretic hormone on cellular conductive pathways in mouse medullary thick ascending limbs of Henle. I. ADH increases transcellular conductance pathways. *J Membr Biol* 80:201–219, 1984
16. HEBERT SC, ANDREOLI TE: Effects of antidiuretic hormone on cellular conductance pathways in mouse medullary thick ascending limbs of Henle. II. Determinants of the ADH-mediated increases in transepithelial voltage and in net Cl^- absorption. *J Membr Biol* 80:221–233, 1984
17. HEBERT SC, ANDREOLI TE: Ionic conductance pathways in the mouse medullary thick ascending limb of Henle. The paracellular pathway and electrogenic Cl^- absorption. *J Gen Physiol* 87:567–590, 1986
18. SCHLATTER E, GREGER R: cAMP increases the basolateral Cl^- conductance in the isolated perfused medullary thick ascending limb of Henle's loop of the mouse. *Pflügers Arch* 405:367–376, 1985
19. MURER H, GREGER R: Membrane transport in the proximal tubule and thick ascending limb of Henle's loop: Mechanisms and their alterations. *Klin Wochenschr* 60:1103–1113, 1982
20. OBERLEITHNER H, GUGGINO W, GIEBISCH G: Mechanism of distal tubular chloride transport in *Amphiuma* kidney. *Am J Physiol* 242:F331–F339, 1982
21. NISHIMURA H, IMAI M, OGAWA M: Sodium chloride and water transport in the renal distal tubule of the rainbow trout. *Am J Physiol* 244:F247–F254, 1983
22. BURG MB: Thick ascending limb of Henle's loop. *Kidney Int* 22:454–464, 1982
23. HEBERT SC, ANDREOLI TE: Control of NaCl transport in the thick ascending limb. *Am J Physiol* 246:F745–F756, 1984
24. KOENIG B, KINNE R: Sodium transport by plasma membranes isolated from cells of the thick ascending limb of Henle's loop. *Fed Proc* 41:1007, 1982
25. HANNAFIN J, KINNE-SAFFRAN E, KINNE R: Sodium and potassium dependent chloride transport in plasma membrane vesicles from mammalian thick ascending limb of Henle's loop and elasmobranch rectal gland. (abstract) *Am Soc Nephrol*, 16th Ann. Meeting, p. 164a, 1984
26. KOENIG B, RICAPITO S, KINNE R: Chloride transport in the thick ascending limb of Henle's loop: Potassium dependence and stoichiometry of the NaCl cotransport system in plasma membrane vesicles. *Pflügers Arch* 399:173–179, 1983
27. EVELOFF J, KINNE R: Sodium-chloride transport in the medullary thick ascending limb of Henle's loop: Evidence for a sodium-chloride cotransport system in plasma membrane vesicles. *J Membr Biol* 72:173–181, 1983
28. ALVO M, CALAMIA J, EVELOFF J: Lack of potassium effect on Na-Cl cotransport in the medullary thick ascending limb. *Am J Physiol* 249:F34–F39, 1985
29. EVELOFF J, FONG J, CALAMIA J: Effect of hyperosmolality and phorbol esters on cation fluxes in medullary thick ascending limb cells. (abstract) *Am Soc Nephrol*, 18th Ann. Meeting, p. 246a, 1985
30. FORBUSH F, III, PALFREY HC: [^3H]Bumetanide binding to membranes isolated from dog kidney outer medulla. *J Biol Chem* 258:11787–11792, 1983
31. OBERLEITHNER H, LANG F, GREGER R, WANG W, GIEBISCH G: Effect of luminal potassium on cellular sodium activity in the early distal tubule of *Amphiuma* kidney. *Pflügers Arch* 396:34–40, 1983
32. OBERLEITHNER H, LANG F, WANG W, GIEBISCH G: Effects of inhibition of chloride transport on intracellular sodium activity in distal amphibian nephron. *Pflügers Arch* 394:55–60, 1982
33. GREGER R, SCHLATTER E: Properties of the basolateral membrane of the cortical thick ascending limb of Henle's loop of rabbit kidney—a model for secondary active chloride transport. *Pflügers Arch* 396:325–334, 1983
34. MOLONY DA, REEVES WB, HEBERT SC, ANDREOLI TE: ADH increases apical $\text{Na}^+:\text{K}^+:2\text{Cl}^-$ entry in mouse medullary thick ascending limbs of Henle. *Am J Physiol* (Submitted)
35. GREGER R, OBERLEITHNER H, SCHLATTER E, CASSOLA AC, WEIDTKE C: Chloride activity in cells of isolated perfused cortical thick ascending limbs of rabbit kidney. *Pflügers Arch* 399:20–34, 1983

36. DiBONA DR: Direct visualization of epithelial morphology in the living amphibian urinary bladder. *J Membr Biol Special Issue*, 1978
37. LEICHTWEISS HP, LÜBBERS DW, WEISS CH, BAUMGÄRTL H, RESCHKE W: The oxygen supply of the rat kidney: Measurement of intrarenal pO_2 . *Pflügers Arch*. 309:328-349, 1969
38. ALCORN D, EMSLIE KR, BOSS BD: Selective distal nephron damage during isolated kidney perfusion. *Kidney Int* 19:638-647, 1981
39. BREZIS M, ROSEN S, SILVA P, EPSTEIN FH: Transport activity modifies thick ascending limb damage in the isolated perfused kidney. *Kidney Int* 25:65-72, 1984
40. SCHULTZ SG: Homocellular regulatory mechanisms in sodium transporting epithelia: Avoidance of extinction by "flush through." *Am J Physiol* 241:F579-F590, 1981
41. WIRZ H: The location of antidiuretic action in the mammalian kidney, in *The Neurohypophysis*, edited by HELLER H. New York, Academic Press, 1957, p. 157-182
42. EDWARDS RM, JACKSON BA, DOUSA TP: Protein kinase activity in isolated tubules of rat renal medulla. *Am J Physiol* 238:F269-F278, 1980
43. HALL DA, VARNEY DM: Effect of vasopressin on electrical potential difference and chloride transport in mouse medullary thick ascending limbs of Henle's loop. *J Clin Invest* 66:792-802, 1980
44. SASAKI S, IMAI M: Effects of vasopressin on water and NaCl transport across the in vitro perfused medullary thick ascending limb of Henle's loop of mouse, rat and rabbit kidneys. *Pflügers Arch* 383:215-221, 1980
45. SHIMIZU K, NAKAO A, NONAKA T, OKA H: Identification of vasopressin and determination of its corticomedullary levels in rat kidney tissue. *Kidney Int* 26:785-790, 1984
46. DEROUFFIGNAC C, CORMAN B, ROINELL N: Stimulation by antidiuretic hormone of electrolyte tubular reabsorption in rat kidney. *Am J Physiol* 244:F156-F164, 1983
47. WORK J, GALLA JH, BOOKER BB, SCHAFER JA, LUKE RG: Effect of ADH on chloride reabsorption in the loop of Henle of the Brattleboro rat. *Am J Physiol* 249:F698-F703, 1985
48. DOUSA TP: Effects of renal medullary solutes on vasopressin-sensitive adenylyl cyclase. *Am J Physiol* 222:657-662, 1972
49. ROY C, LEBARS NC, JARD S: Vasopressin-sensitive kidney adenylyl cyclase. Differential effects of monovalent ions on stimulations by fluoride, vasopressin and guanylyl 5'-imidodiphosphate. *Eur J Biochem* 78:325-332, 1977
50. DERUBERTIS FR, CRANEN PA: Effect of osmolality and oxygen availability on soluble cyclic AMP-dependent protein kinase activity of rat inner medulla. *J Clin Invest* 62:1210-1221, 1978
51. EDWARDS RM, JACKSON BA, DOUSA TP: ADH-sensitive cAMP system in papillary collecting duct: Effects of osmolality and PGE. *Am J Physiol* 240:F311-F318, 1981
52. TORIKAI S, IMAI M: Effects of solute concentration on vasopressin stimulated cyclic AMP generation in the rat medullary thick ascending limbs of Henle's loop. *Pflügers Arch* 400:306-308, 1984
53. BECK TR, DUNN MJ: The relationship of antidiuretic hormone and renal prostaglandins. *Miner Electrol Metab* 6:46-59, 1981
54. DERUBERTIS FR, CRAVEN PA: Prostaglandin metabolism in the kidney: Physiologic implications. *Sem Nephrol* 2:302-315, 1982
55. STOKES JB: Integrated actions of renal medullary prostaglandins in the control of water excretion. *Am J Physiol* 240:F471-F480, 1981
56. CULPEPPER RM, ANDREOLI TE: Interactions among prostaglandin E_2 , antidiuretic hormone, and cyclic adenosine monophosphate in modulating Cl^- absorption in single mouse medullary thick ascending limbs of Henle. *J Clin Invest* 71:1588-1601, 1983
57. FINE LG, TRIZNA W: Influence of prostaglandins on sodium transport of isolated medullary nephron segments. *Am J Physiol* 232:F383-F390, 1977
58. KUAKER ML: Prostaglandin E_2 effect from the luminal side on renal tubular ^{22}Na efflux: Tracer microinjection studies. *Proc Soc Exp Biol Med* 154:274-277, 1977
59. GANGULI M, TOBIAN L, AZAR S, O'DONNELL M: Evidence that prostaglandin synthesis inhibitors increase the concentration of sodium and chloride in rat renal medulla. *Circ Res (suppl)* I:1139, 1977
60. HIGASHIHARA E, STOKES JB, KOKKO JP, CAMPBELL WB, DuBOSE TD: Cortical and papillary micropuncture examination of chloride transport in segments of the rat kidney during inhibition of prostaglandin production: Possible role for prostaglandins in chloruresis of acute volume expansion. *J Clin Invest* 64:1277-1287, 1979
61. KIRCHNER KA: Prostaglandin inhibitors alter loop segment chloride uptake during furosemide diuresis. *Am J Physiol* 248:F698-F704, 1985
62. STOKES JB: Effect of prostaglandin E_2 on chloride transport across the rabbit thick ascending limb of Henle. *J Clin Invest* 64:495-502, 1979
63. CULPEPPER RM, ANDREOLI TE: PGE₂, forskolin, and cholera toxin interactions in modulating NaCl transport in mouse mTALH. *Am J Physiol* 247:F784-F792, 1984
64. TORIKAI S, KUROKAWA K: Effect of PGE₂ on vasopressin-dependent cell cAMP in isolated nephron segments. *Am J Physiol* 245:F58-F66, 1983
65. NADLER SP, HEBERT SC, BRENNER BM: PGE₂, forskolin, and cholera toxin interactions in the rabbit cortical collecting tubule. *Am J Physiol* (In press)
66. SMITH WL, WILKIN GP: Immunocytochemistry of prostaglandin endoperoxide-forming cyclooxygenases: The detection of the cyclooxygenases in rat, rabbit and guinea pig kidneys by immunofluorescence. *Prostaglandins* 13:873-892, 1977
67. CRAVEN PA, DERUBERTIS FR: Effects of vasopressin and urea on Ca^{2+} -calmodulin-dependent renal prostaglandin E. *Am J Physiol* 241:F649-F658, 1981
68. ZIPSER RD, LITTLE TW, WILSON W, DUKE R: Dual effects of antidiuretic hormone on urinary prostaglandin E_2 excretion in man. *J Clin Endocrinol Metab* 53:522-526, 1981
69. CAMPBELL HT, CRAVEN PA, DERUBERTIS FR: Suppression of basal and vasopressin-responsive urinary prostaglandin E excretion by water deprivation. (abstract) *Clin Res* 30:444A, 1982
70. KIRCHENBAUM MA, SERROS ER: Effects of alterations in urinary flow rate on prostaglandin E excretion in conscious dogs. *Am J Physiol* 238:F107-F111, 1980
71. HEBERT SC, CULPEPPER RM, ANDREOLI TE: NaCl transport in mouse medullary thick ascending limbs. III. Modulation of the ADH effect by peritubular osmolality. *Am J Physiol* 241:F443-F451, 1981
72. LEVI J, COBURN J, KLEEMAN CR: Mechanism of the antidiuretic effect of α -adrenergic stimulation in man. *Arch Int Med* 136:25-29, 1976
73. KLEIN LA, LIEBERMAN B, LAKS M, KLEEMAN CR: Interrelated effects of antidiuretic hormone and adrenergic drugs on water metabolism. *Am J Physiol* 221:1657-1665, 1971
74. LEVI J, GRINBLAT J, KLEEMAN CR: Effect of isoproterenol on water diuresis in the rat with congenital diabetes insipidus. *Am J Physiol* 22:1728-1732, 1971
75. POLHEMUS RE, HALL DA: Effect of catecholamines on the potential difference and chloride efflux in the mouse thick ascending limb of Henle's loop. (abstract) *Kidney Int* 19:253, 1981
76. CULPEPPER RM: Multiple regulators of adenylyl cyclase (AC) in mouse medullary thick ascending limbs (mTALH): Significance for NaCl transport. (abstract) *Clin Res* 32:65A, 1984
77. MOREL F, IMBERT-TEBOUL M, LEBOUFFANT F, HUS-CITHAREL A, MONTGUT M: Multiple hormonal control of adenylyl cyclase in distal segments of the rat kidney. *Kidney Int* 21(suppl 11):55-62, 1982
78. BARAJAS L, WANG P, POWERS K, NISHIO S: Identification of renal neuroeffector junctions by electron microscopy of re-embedded light microscopic autoradiograms of semithin sections. *J Ultrastruct Res* 77:379-385, 1981
79. BARAJAS L, POWERS K, WANG P: Innervation of the renal cortical tubules: A quantitative study. *Am J Physiol* 247:F50-F60, 1984
80. DiBONA GF, SAWIN LL: Effect of renal nerve stimulation on NaCl and H_2O transport in Henle's loop of the rat. *Am J Physiol* 243:F576-F580, 1982
81. BENCÁTH P, SZÉNÁSI G, TAKÁCS L: Water and electrolyte transport in Henle's loop and distal tubule after renal sympathectomy.

- tomy in the rat. *Am J Physiol* 249:F308-F314, 1985
82. BATTILANA CA, DOBYAN DC, LACY FB, BHATTACHARYA J, JOHNSTON PA, JAMISON RL: Effect of chronic potassium loading on potassium secretion by the pars recta or descending limb of the juxtamedullary nephron in the rat. *J Clin Invest* 62:1093-1103, 1978
 83. JAMISON RL, WORK J, SCHAFER JA: New pathways for potassium transport in the kidney. *Am J Physiol* 242:F297-F312, 1982
 84. STOKES JB: Consequences of potassium recycling in the renal medulla. Effects on ion transport by the medullary thick ascending limb of Henle's loop. *J Clin Invest* 70:219-229, 1982
 85. ELALOUF J-M, ROINEL N, DEROUFFIGNAC C: Effects of dDAVP on rat juxtamedullary nephrons: Stimulation of medullary K recycling. *Am J Physiol* 249:F291-F298, 1985
 86. KIRCHNER KA: Effect of acute potassium infusion on loop segment chloride reabsorption in the rat. *Am J Physiol* 244:F599-F605, 1983
 87. LUKE RG, BOOKER BB, GALLA JH: Effect of potassium depletion on chloride transport in the loop of Henle in the rat. *Am J Physiol* 248:F682-F687, 1985
 88. KIM JK, JACKSON BA, EDWARDS RM, DOUSA TP: Effect of potassium depletion on the vasopressin-sensitive cyclic AMP system in rat outer medullary tubules. *J Lab Clin Med* 99:29-38, 1982
 89. MANITIUS A, LEVITIN H, BECK D, EPSTEIN FH: On the mechanism of impairment of renal concentrating ability in hypercalcemia. *J Clin Invest* 39:693-697, 1960
 90. LEVI M, PETERSON L, BERL T: Mechanism of concentrating defect in hypercalcemia. Role of polydipsia nad prostaglandins. *Kidney Int* 23:489-497, 1983
 91. KUSANO E, MURAYAMA N, WERNES JL, CHRISTENSEN S, HOMMA S, YUSUFI ANK, DOUSA TP: Effects of calcium on the vasopressin-sensitive cAMP metabolism in medullary tubules. *Am J Physiol* 249:F956-F966, 1985
 92. SAIKIA TC: Composition of the renal cortex and medulla of rats during water diuresis and antidiuresis. *Q J Exp Physiol* 50:146157, 1965
 93. HAI MA, THOMAS S: The time-course of changes in renal tissue composition during lysine vasopressin infusion in the rat. *Pflügers Arch* 310:297-319, 1969
 94. SCHMIDT-NIELSEN B, GRAVES B, ROTH J: Water removal and solute additions determining increases in renal medullary osmolality. *Am J Physiol* 244:F472-F482, 1983
 95. WANGEMANN P, WITTNER M, DiSTEFANO A, ENGLERT HC, LANG HJ, SCHLATTER E, GREGER R: Cl^- -channel blockers in the thick ascending limb of the loop of Henle. Structure activity relationship. *Pflügers Arch* (submitted)
 96. AL-ZAHID G, SCHAFER JA, TROUTMAN SL, ANDREOLI TE: Effects of antidiuretic hormone on water and solute permeation and the activation energies for these processes in mammalian cortical collecting tubules: Evidence for parallel ADH-sensitive pathways for water and solute diffusion in luminal plasma membranes. *J Membr Biol* 31:103-129, 1977
 97. HAYS RM, FRANKI N: The role of water diffusion in the action of vasopressin. *J Membr Biol* 2:263-276, 1970
 98. HEBERT SC, ANDREOLI TE: Interactions of temperature and ADH on transport processes in cortical collecting tubules. *Am J Physiol* 238:F470-F480, 1980
 99. LEVINE SD, JACOBY M, FINKELSTEIN A: The water permeability of toad urinary bladder. II. The value of $P_f/P_d(w)$ for the antidiuretic hormone-induced water permeation pathway. *J Gen Physiol* 83:543-561, 1984
 100. LI H-Y S, PALMER LG, EDELMAN IS, LINDEMANN B: The role of sodium-channel density in the natriuretic response of the toad urinary bladder to an antidiuretic hormone. *J Membr Biol* 64:77-89, 1982
 101. SCHAFER JA, TROUTMAN SL: Vasopressin stimulates Rubidium secretion in the isolated rat cortical collecting tubule. (abstract) *Kidney Int* 27:320A, 1985
 102. HAYS RM, BOURGUET J, CHEVALIER J: Membrane fusion in the action of ADH, determined with an ultrarapid freezing technique. (abstract) *Proc Am Soc Nephrol* p. 149A, 1981